

# VITALIZE<sup>®</sup>



## VITALIZE<sup>®</sup> EQUINE RECOVERY GEL

### Examples of When and How to Use



#### SHOWING

Administer 1 tube approximately 2 hours prior to showing. For overly stressed or anxious horses, administer 1 tube the night before competition as well as 2 hours prior.



#### HAULING

Administer 1 tube approximately 2 hours prior to hauling. For long trips, administer 1 additional tube every 4 hours while hauling up to two times a day.



#### WEANING

Administer 15 mL to foal for 2 days prior to weaning, on the day of weaning, and for 2 days following weaning.



#### DIGESTIVE UPSET

Administer 1 tube as soon as digestive upset is noticed and every 4 hours afterward until condition is improved up to two times a day.



#### DIARRHEA

Administer 1 tube when diarrhea is noticed.



#### NOT EATING OR DRINKING

Administer 1 tube immediately and then every 4 hours afterward until condition is improved up to two times a day.



#### ON ANTIBIOTICS

Administer 1 tube twice daily during course of antibiotics and for 1 week afterward.



#### OTHER STRESS (WEATHER CHANGES, SOCIAL STRESS, ETC.)

Administer 1 tube every 4 to 24 hours depending on extent of stress up to two times a day.



A *good gut* FEELING



# VITALIZE<sup>®</sup>



## VITALIZE<sup>®</sup> DOG RECOVERY GEL

### Examples of When and How to Use



#### TRAVELING

Administer 2 mL per 10 lbs. of body weight before traveling begins and every 4 hours afterward while traveling.



#### COMPETING OR PERFORMANCE

Administer 2 mL per 10 lbs. of body weight approximately 2 hours before competing. For overly stressed or anxious dogs, administer the same dose the night before as well as 2 hours before.



#### VOMITING OR DIARRHEA

Administer 2 mL per 10 lbs. of body weight when condition is noticed.



#### BLOATING

Administer 2 mL per 10 lbs. of body weight when condition is noticed.



#### ALLERGIES OR SKIN PROBLEMS

Administer 2 mL per 10 lbs. of body weight every 4 to 24 hours depending on severity.



#### TOXICITY OR POISONING

Administer 2 mL per 10 lbs. of body weight as soon as noticed.



#### ON ANTIBIOTICS

Administer 2 mL per 10 lbs. of body weight twice daily during course of antibiotics and for 1 week afterward.



#### OTHER STRESS (STORMS, INJURY, ETC.)

Administer 2 mL per 10 lbs. of body weight every 4 to 24 hours depending on extent of stress.



A *good gut* FEELING